


Arachidonic acid metabolism



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://creativecommons.org/licenses/by/4.0/).

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)


Reactome database release: 74

This document contains 8 pathways and 3 reactions ([see Table of Contents](#))

Arachidonic acid metabolism ↗

Stable identifier: R-CEL-2142753

Inferred from: Arachidonic acid metabolism (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

AWAT1 transfers acyl group from acyl-CoA to ARACOH, forming wax esters ↗


Location: [Arachidonic acid metabolism](#)

Stable identifier: R-CEL-5696424

Type: transition

Compartments: endoplasmic reticulum membrane, endoplasmic reticulum lumen

Inferred from: [AWAT1 transfers acyl group from acyl-CoA to ARACOH, forming wax esters \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

FAAH hydrolyses AEA to AA and ETA ↗


Location: [Arachidonic acid metabolism](#)

Stable identifier: R-CEL-5693742

Type: transition

Compartments: endoplasmic reticulum membrane, cytosol

Inferred from: [FAAH hydrolyses AEA to AA and ETA \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

FAAH2 hydrolyses AEA to AA and ETA [↗](#)


Location: [Arachidonic acid metabolism](#)

Stable identifier: R-CEL-5693751

Type: transition

Compartments: lipid droplet

Inferred from: [FAAH2 hydrolyses AEA to AA and ETA \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Synthesis of Prostaglandins (PG) and Thromboxanes (TX) ↗

Location: Arachidonic acid metabolism

Stable identifier: R-CEL-2162123

Inferred from: Synthesis of Prostaglandins (PG) and Thromboxanes (TX) (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


More details and caveats of the event inference in Reactome. For details on PANTHER see also: http://www.pantherdb.org/about.jsp

Synthesis of Leukotrienes (LT) and Eoxins (EX) ↗

Location: Arachidonic acid metabolism

Stable identifier: R-CEL-2142691

Inferred from: Synthesis of Leukotrienes (LT) and Eoxins (EX) (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Synthesis of 5-icosatetraenoic acids ↗

Location: Arachidonic acid metabolism

Stable identifier: R-CEL-2142688

Inferred from: Synthesis of 5-icosatetraenoic acids (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Synthesis of 15-eicosatetraenoic acid derivatives ↗

Location: Arachidonic acid metabolism

Stable identifier: R-CEL-2142770

Inferred from: Synthesis of 15-eicosatetraenoic acid derivatives (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Synthesis of 12-eicosatetraenoic acid derivatives ↗

Location: Arachidonic acid metabolism

Stable identifier: R-CEL-2142712

Inferred from: Synthesis of 12-eicosatetraenoic acid derivatives (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) ↗

Location: Arachidonic acid metabolism

Stable identifier: R-CEL-2142816

Inferred from: Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE) (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Synthesis of epoxy (EET) and dihydroxyeicosatrienoic acids (DHET) ↗

Location: Arachidonic acid metabolism

Stable identifier: R-CEL-2142670

Inferred from: Synthesis of epoxy (EET) and dihydroxyeicosatrienoic acids (DHET) (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Table of Contents

Introduction	1
☒ Arachidonic acid metabolism	2
↳ AWAT1 transfers acyl group from acyl-CoA to ARACOH, forming wax esters	3
↳ FAAH hydrolyses AEA to AA and ETA	4
↳ FAAH2 hydrolyses AEA to AA and ETA	5
☒ Synthesis of Prostaglandins (PG) and Thromboxanes (TX)	6
☒ Synthesis of Leukotrienes (LT) and Eoxins (EX)	7
☒ Synthesis of 5-eicosatetraenoic acids	8
☒ Synthesis of 15-eicosatetraenoic acid derivatives	9
☒ Synthesis of 12-eicosatetraenoic acid derivatives	10
☒ Synthesis of (16-20)-hydroxyeicosatetraenoic acids (HETE)	11
☒ Synthesis of epoxy (EET) and dihydroxyeicosatrienoic acids (DHET)	12
Table of Contents	13