


Regulation of lipid metabolism by Peroxisome proliferator-activated receptor alpha (PPARalpha)



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://creativecommons.org/licenses/by/4.0/).

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 70


This document contains 1 pathway and 3 reactions ([see Table of Contents](#))

Regulation of lipid metabolism by Peroxisome proliferator-activated receptor alpha (PPARalpha) ↗

Stable identifier: R-DRE-400206

Compartments: nucleoplasm, cytosol

Inferred from: Regulation of lipid metabolism by Peroxisome proliferator-activated receptor alpha (PPARalpha) (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

PPARA binds RXRA forming PPARA:RXRA heterodimer at promoter of target gene ↗


Location: Regulation of lipid metabolism by Peroxisome proliferator-activated receptor alpha (PPARα)

Stable identifier: R-DRE-400204

Type: binding

Compartments: nucleoplasm

Inferred from: PPARA binds RXRA forming PPARA:RXRA heterodimer at promoter of target gene (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: PPARA:RXRA heterodimer recruits corepressors

PPARA:RXRA heterodimer recruits corepressors ↗


Location: Regulation of lipid metabolism by Peroxisome proliferator-activated receptor alpha (PPARalpha)

Stable identifier: R-DRE-400183

Type: binding

Compartments: nucleoplasm

Inferred from: PPARA:RXRA heterodimer recruits corepressors (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: PPARA binds RXRA forming PPARA:RXRA heterodimer at promoter of target gene

Followed by: Fatty acid ligands activate PPARA

Fatty acid ligands activate PPARA ↗


Location: Regulation of lipid metabolism by Peroxisome proliferator-activated receptor alpha (PPARalpha)

Stable identifier: R-DRE-400143

Type: transition

Compartments: nucleoplasm

Inferred from: Fatty acid ligands activate PPARA (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [PPARA:RXRA heterodimer recruits corepressors](#)

Table of Contents

Introduction	1
⚡ Regulation of lipid metabolism by Peroxisome proliferator-activated receptor alpha (PPARalpha)	2
➤ PPARA binds RXRA forming PPARA:RXRA heterodimer at promoter of target gene	3
➤ PPARA:RXRA heterodimer recruits corepressors	4
➤ Fatty acid ligands activate PPARA	5
Table of Contents	6