


# GLI3 is processed to GLI3R by the protea- some



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](#).

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)


Reactome database release: 70

This document contains 1 pathway and 5 reactions ([see Table of Contents](#))

## GLI3 is processed to GLI3R by the proteasome ↗

**Stable identifier:** R-DRE-5610785

**Inferred from:** GLI3 is processed to GLI3R by the proteasome (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

## PKA phosphorylates GLI3 ↗


**Location:** GLI3 is processed to GLI3R by the proteasome

**Stable identifier:** R-DRE-5610720

**Type:** transition

**Compartments:** ciliary base

**Inferred from:** PKA phosphorylates GLI3 (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

**Followed by:** CK1 phosphorylates p-GLI3

## CK1 phosphorylates p-GLI3 ↗


**Location:** GLI3 is processed to GLI3R by the proteasome

**Stable identifier:** R-DRE-5610722

**Type:** transition

**Compartments:** cytosol

**Inferred from:** CK1 phosphorylates p-GLI3 (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

**Preceded by:** PKA phosphorylates GLI3

**Followed by:** GSK3 phosphorylates p-GLI3

## GSK3 phosphorylates p-GLI3 ↗


**Location:** GLI3 is processed to GLI3R by the proteasome

**Stable identifier:** R-DRE-5610732

**Type:** transition

**Compartments:** cytosol

**Inferred from:** GSK3 phosphorylates p-GLI3 (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

**Preceded by:** CK1 phosphorylates p-GLI3

**Followed by:** SCF(beta-TrCP) ubiquitinates p-GLI3

## SCF(beta-TrCP) ubiquitinates p-GLI3 [↗](#)


**Location:** GLI3 is processed to GLI3R by the proteasome

**Stable identifier:** R-DRE-5610746

**Type:** transition

**Compartments:** cytosol

**Inferred from:** SCF(beta-TrCP) ubiquitinates p-GLI3 (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

**Preceded by:** GSK3 phosphorylates p-GLI3

**Followed by:** GLI3 is partially degraded by the proteasome to yield the GLI3 repressor

## GLI3 is partially degraded by the proteasome to yield the GLI3 repressor ↗


**Location:** GLI3 is processed to GLI3R by the proteasome

**Stable identifier:** R-DRE-5610754

**Type:** omitted

**Compartments:** cytosol

**Inferred from:** GLI3 is partially degraded by the proteasome to yield the GLI3 repressor (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

**Preceded by:** SCF(beta-TrCP) ubiquitinates p-GLI3

# Table of Contents

Introduction	1
☒ GLI3 is processed to GLI3R by the proteasome	2
↳ PKA phosphorylates GLI3	3
↳ CK1 phosphorylates p-GLI3	4
↳ GSK3 phosphorylates p-GLI3	5
↳ SCF(beta-TrCP) ubiquitinates p-GLI3	6
☒ GLI3 is partially degraded by the proteasome to yield the GLI3 repressor	7
Table of Contents	8