


# Sema4D induced cell migration and growth-cone collapse



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the [Reactome Textbook](#).

02/10/2022

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 82


This document contains 1 pathway and 5 reactions ([see Table of Contents](#))

## Sema4D induced cell migration and growth-cone collapse ↗

**Stable identifier:** R-GGA-416572

**Compartments:** plasma membrane

**Inferred from:** [Sema4D induced cell migration and growth-cone collapse \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

## SEMA4D interacts with Plexin-B1:ErbB2 ↗


**Location:** [Sema4D induced cell migration and growth-cone collapse](#)

**Stable identifier:** R-GGA-373750

**Type:** transition

**Compartments:** plasma membrane, cytosol

**Inferred from:** [SEMA4D interacts with Plexin-B1:ErbB2 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

**Followed by:** [LARG and PDZ-RhoGEF binds to Plexin-B1](#)

## LARG and PDZ-RhoGEF binds to Plexin-B1 [↗](#)


**Location:** [Sema4D induced cell migration and growth-cone collapse](#)

**Stable identifier:** R-GGA-416594

**Type:** binding

**Compartments:** plasma membrane, cytosol

**Inferred from:** [LARG and PDZ-RhoGEF binds to Plexin-B1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

**Preceded by:** [SEMA4D interacts with Plexin-B1:ErbB2](#)

**Followed by:** [Activation of Rho by LARG and PDZ-RhoGEF](#)

## Activation of Rho by LARG and PDZ-RhoGEF ↗


**Location:** [Sema4D induced cell migration and growth-cone collapse](#)

**Stable identifier:** R-GGA-416588

**Type:** transition

**Compartments:** plasma membrane, cytosol

**Inferred from:** [Activation of Rho by LARG and PDZ-RhoGEF \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

**Preceded by:** [LARG and PDZ-RhoGEF binds to Plexin-B1](#)

**Followed by:** [ROCK activation by RHO](#)

## ROCK activation by RHO ↗


**Location:** [Sema4D induced cell migration and growth-cone collapse](#)

**Stable identifier:** R-GGA-419049

**Type:** binding

**Compartments:** plasma membrane, cytosol

**Inferred from:** [ROCK activation by RHO \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

**Preceded by:** [Activation of Rho by LARG and PDZ-RhoGEF](#)

## ROCK1,2 bind ROCKi ↗


**Location:** [Sema4D induced cell migration and growth-cone collapse](#)

**Stable identifier:** R-GGA-9680443

**Type:** binding

**Compartments:** cytosol

**Inferred from:** [ROCK1,2 bind ROCKi \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

# Table of Contents

Introduction	1
☒ Sema4D induced cell migration and growth-cone collapse	2
↳ SEMA4D interacts with Plexin-B1:ErbB2	3
↳ LARG and PDZ-RhoGEF binds to Plexin-B1	4
↳ Activation of Rho by LARG and PDZ-RhoGEF	5
↳ ROCK activation by RHO	6
↳ ROCK1,2 bind ROCKi	7
Table of Contents	8