

Association of LBP with LPS

Gay, NJ., Gillespie, ME., Granucci, F., Shamovsky, V., Zanoni, I., de Bono, B.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 75


This document contains 1 reaction ([see Table of Contents](#))

Association of LBP with LPS [↗](#)

Stable identifier: R-HSA-166015

Type: binding

Compartments: extracellular region



Lipopolysaccharide-binding protein (LBP) is a ~60-kDa serum glycoprotein which transfers LPS to both membrane-bound and soluble CD14. The LPS binding site of LBP consists of basic residues that bind the phosphorylated head of the bacterial lipid A.

LBP is an acute-phase opsonin that binds gram-negative bacteria and bacterial fragments and promote the interaction of coated bacteria with phagocytes.

Literature references

Wright, SD., Tobias, PS., Ulevitch, RJ., Ramos, RA. (1989). Lipopolysaccharide (LPS) binding protein opsonizes LPS-bearing particles. *J Exp Med*, 170, 1231-41. [↗](#)

Editions

2005-08-16	Authored	de Bono, B.
2006-04-24	Reviewed	Gay, NJ.
2010-11-16	Edited	Shamovsky, V.
2010-11-30	Reviewed	Gillespie, ME.
2012-11-13	Reviewed	Zanoni, I., Granucci, F.