

Interaction of integrin alphaIIb beta3 with von Willebrand factor

Garapati, P V., Geiger, B., Horwitz, AR., Humphries, MJ., Hynes, R., Yamada, KM.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 70


This document contains 1 reaction ([see Table of Contents](#))

Interaction of integrin alphaIIbeta3 with von Willebrand factor ↗

Stable identifier: R-HSA-216072

Type: binding

Compartments: extracellular region, plasma membrane



Integrin alphaIIbeta3 (glycoprotein IIb-IIIa, GP IIb-IIIa) is one of the major platelet receptor that is involved in aggregation and adhesion of platelets. In resting stage alphaIIbeta3 is inactive and does not interact with its ligands but upon activation or vascular injury it binds to the ECM protein von Willebrand factor (vWf) and stimulate the platelet aggregation.

Literature references

Giuliano, S., Nesbitt, WS., Rooney, M., Jackson, SP. (2003). Bidirectional integrin alphaIIbeta3 signalling regulating platelet adhesion under flow: contribution of protein kinase C. *Biochem J*, 372, 163-72. ↗

Nesbitt, WS., Kulkarni, S., Giuliano, S., Goncalves, I., Dopheide, SM., Yap, CL. et al. (2002). Distinct glycoprotein Ib/V/IX and integrin alpha IIbeta 3-dependent calcium signals cooperatively regulate platelet adhesion under flow. *J Biol Chem*, 277, 2965-72. ↗

Editions

2008-03-11	Edited	Garapati, P V.
2008-05-07	Authored	Geiger, B., Horwitz, AR.
2008-05-07	Reviewed	Humphries, MJ., Yamada, KM., Hynes, R.