

# AP-1 stimulates IL1A transcription

Orlic-Milacic, M., Samarajiwa, S.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 70


This document contains 1 reaction ([see Table of Contents](#))

## AP-1 stimulates IL1A transcription ↗

**Stable identifier:** R-HSA-4568740

**Type:** omitted

**Compartments:** nucleoplasm, extracellular region



The AP-1 complex (FOS:JUN) binds IL1A promoter and stimulates IL1A transcription (Bailly et al. 1996 Alheim et al. 1996, Burleson et al. 1996, Huang et al. 1999, Niu et al. 2004, Moerman-Herzog and Barger 2012, Murakami et al. 2013).

### Literature references

- Alheim, K., McDowell, TL., Symons, JA., Duff, GW., Bartfai, T. (1996). An AP-1 site is involved in the NGF induction of IL-1 alpha in PC12 cells. *Neurochem. Int.*, 29, 487-96. ↗
- Bailly, S., Fay, M., Israël, N., Gougerot-Pocidallo, MA. (1996). The transcription factor AP-1 binds to the human interleukin 1 alpha promoter. *Eur. Cytokine Netw.*, 7, 125-8. ↗
- Huang, TT., Vinci, JM., Lan, L., Jeffrey, JJ., Wilcox, BD. (1999). Serotonin-inducible transcription of interleukin-1alpha in uterine smooth muscle cells requires an AP-1 site: cloning and partial characterization of the rat IL-1alpha promoter. *Mol. Cell. Endocrinol.*, 152, 21-35. ↗
- Moerman-Herzog, AM., Barger, SW. (2012). A polymorphism in the upstream regulatory region of the interleukin-1? gene confers differential binding by transcription factors of the AP-1 family. *Life Sci.*, 90, 975-9. ↗
- Niu, J., Li, Z., Peng, B., Chiao, PJ. (2004). Identification of an autoregulatory feedback pathway involving interleukin-1alpha in induction of constitutive NF-kappaB activation in pancreatic cancer cells. *J. Biol. Chem.*, 279, 16452-62. ↗

### Editions

2013-09-03	Reviewed	Samarajiwa, S.
2013-09-16	Authored, Edited	Orlic-Milacic, M.