

# RUNX1 binds the SWI/SNF complex

Chuang, L.S., Ito, Y., Orlic-Milacic, M.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 74


This document contains 1 reaction ([see Table of Contents](#))

## RUNX1 binds the SWI/SNF complex [↗](#)

**Stable identifier:** R-HSA-8938217

**Type:** binding

**Compartments:** nucleoplasm



The RUNX1:CBFB complex can bind the SWI/SNF chromatin remodeling complex by directly interacting with at least three subunits of the SWI/SNF: BRG1 (SMARCA4), BAF155 (SMARCC1) and INI1 (SMARCB1). RUNX1 recruits the SWI/SNF complex to many of its target promoters, which is associated with changes in histone modifications at these promoters, but precise changes and their effect on gene expression have not been fully elucidated (Bakshi et al. 2010).

### Literature references

Bakshi, R., Hassan, MQ., Pratap, J., Lian, JB., Montecino, MA., van Wijnen, AJ. et al. (2010). The human SWI/SNF complex associates with RUNX1 to control transcription of hematopoietic target genes. *J. Cell. Physiol.*, 225, 569-76. [↗](#)

### Editions

2016-09-14	Authored	Orlic-Milacic, M.
2016-12-20	Reviewed	Ito, Y., Chuang, LS.
2017-05-09	Edited	Orlic-Milacic, M.