


Resolution of Sister Chromatid Cohesion



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://creativecommons.org/licenses/by/4.0/).

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 70

This document contains 1 pathway and 8 reactions ([see Table of Contents](#))

CDK1 phosphorylates CDCA5 (Sororin) at chromosomal arms ↗


Location: [Resolution of Sister Chromatid Cohesion](#)

Stable identifier: R-MMU-2468293

Type: transition

Compartments: chromosome, cytosol

Inferred from: [CDK1 phosphorylates CDCA5 \(Sororin\) at chromosomal arms \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: [Phosphorylation of cohesin by PLK1 at chromosomal arms](#)

Phosphorylation of cohesin by PLK1 at chromosomal arms ↗


Location: [Resolution of Sister Chromatid Cohesion](#)

Stable identifier: R-MMU-2466068

Type: transition

Compartments: chromosome, cytosol

Inferred from: [Phosphorylation of cohesin by PLK1 at chromosomal arms \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [CDK1 phosphorylates CDCA5 \(Sororin\) at chromosomal arms](#)

Followed by: [Resolution of sister chromatids](#)

Resolution of sister chromatids ↗


Location: [Resolution of Sister Chromatid Cohesion](#)

Stable identifier: R-MMU-2467794

Type: dissociation

Compartments: chromosome, cytosol

Inferred from: [Resolution of sister chromatids \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [Phosphorylation of cohesin by PLK1 at chromosomal arms](#)

Followed by: [Deacetylation of cohesin](#)

Deacetylation of cohesin ↗


Location: [Resolution of Sister Chromatid Cohesion](#)

Stable identifier: R-MMU-2545253

Type: transition

Compartments: cytosol

Inferred from: [Deacetylation of cohesin \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [Resolution of sister chromatids](#)

Kinetochores assembly ↗


Location: [Resolution of Sister Chromatid Cohesion](#)

Stable identifier: R-MMU-2484822

Type: omitted

Compartments: chromosome, centromeric region, cytosol

Inferred from: [Kinetochores assembly \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: [CDK1 phosphorylates CDCA5 \(Sororin\) at centromeres](#)

CDK1 phosphorylates CDCA5 (Sororin) at centromeres ↗


Location: [Resolution of Sister Chromatid Cohesion](#)

Stable identifier: R-MMU-2468287

Type: transition

Compartments: chromosome, centromeric region, cytosol

Inferred from: [CDK1 phosphorylates CDCA5 \(Sororin\) at centromeres \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [Kinetochores assembly](#)

Followed by: [Phosphorylation of cohesin by PLK1 at centromeres](#)

Phosphorylation of cohesin by PLK1 at centromeres ↗


Location: [Resolution of Sister Chromatid Cohesion](#)

Stable identifier: R-MMU-1638803

Type: transition

Compartments: chromosome, centromeric region, cytosol

Inferred from: [Phosphorylation of cohesin by PLK1 at centromeres \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [CDK1 phosphorylates CDCA5 \(Sororin\) at centromeres](#)

Followed by: [PP2A-B56 dephosphorylates centromeric cohesin](#)

PP2A-B56 dephosphorylates centromeric cohesin [↗](#)


Location: [Resolution of Sister Chromatid Cohesion](#)

Stable identifier: R-MMU-1638821

Type: transition

Compartments: cytosol, chromosome

Inferred from: [PP2A-B56 dephosphorylates centromeric cohesin \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [Phosphorylation of cohesin by PLK1 at centromeres](#)

Table of Contents

Introduction	1
⚡ Resolution of Sister Chromatid Cohesion	2
↳ CDK1 phosphorylates CDCA5 (Sororin) at chromosomal arms	3
↳ Phosphorylation of cohesin by PLK1 at chromosomal arms	4
↳ Resolution of sister chromatids	5
↳ Deacetylation of cohesin	6
↳ Kinetochores assembly	7
↳ CDK1 phosphorylates CDCA5 (Sororin) at centromeres	8
↳ Phosphorylation of cohesin by PLK1 at centromeres	9
↳ PP2A-B56 dephosphorylates centromeric cohesin	10
Table of Contents	11