

EphBs binds Ptk2

Garapati, P V., Ip, NY.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

31/10/2020

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 74


This document contains 1 reaction ([see Table of Contents](#))

EphBs binds Ptk2 [↗](#)

Stable identifier: R-MMU-3928630

Type: binding

Compartments: plasma membrane, cytosol



Focal adhesion kinase 1 (PTK2, FAK, FAK1) acts downstream of EPHB receptors in hippocampal neurons and the EPHB2-FAK signaling contributes to the dendritic spine morphogenesis and synapse maturation by suppressing the activity of actin severing cofilin through phosphorylation. Activation of EPHBs by ephrin-B (EFNB) stimulates the binding of FAK to EPHB. Knock out of FAK in mature neurons induces a shift of mushroom shaped mature dendritic spines to long filopodia like structures, suggesting that synapse formation or maturation is affected in FAK^{-/-} neurons (Shi et al. 2009, Moeller et al. 2006).

Literature references

Moeller, ML., Shi, Y., Reichardt, LF., Ethell, IM. (2006). EphB receptors regulate dendritic spine morphogenesis through the recruitment/phosphorylation of focal adhesion kinase and RhoA activation. *J. Biol. Chem.*, 281, 1587-98. [↗](#)

Shi, Y., Pontrello, CG., DeFea, KA., Reichardt, LF., Ethell, IM. (2009). Focal adhesion kinase acts downstream of EphB receptors to maintain mature dendritic spines by regulating cofilin activity. *J. Neurosci.*, 29, 8129-42. [↗](#)

Editions

2013-07-23	Authored, Edited	Garapati, P V.
2014-05-19	Reviewed	Ip, NY.