

# Phosphorylation of syk

Garapati, P V., Roncagalli, R.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 70


This document contains 1 reaction ([see Table of Contents](#))

## Phosphorylation of syk ↗

**Stable identifier:** R-RNO-2054095

**Type:** transition

**Compartments:** cytosol



Multiple sites of phosphorylation are known to exist in SYK, which both regulate its activity and also serve as docking sites for other proteins. Some of these sites include Y131 of interdomain A, Y323, Y348, and Y352 of interdomain B, and Y525 and Y526 within the activation loop of the kinase domain and Y630 in the C-terminus (Zhang et al. 2002, Lupher et al. 1998, Furlong et al. 1997). Phosphorylation of these tyrosine residues disrupts autoinhibitory interactions and results in kinase activation even in the absence of phosphorylated ITAM tyrosines (Tsang et al. 2008). SYK is primarily phosphorylated by Src family kinases and this acts as an initiating trigger by generating few molecules of activated SYK which are then involved in major SYK autophosphorylation (Hillal et al. 1997).

### Literature references

Zhang, J., Billingsley, ML., Kincaid, RL., Siraganian, RP. (2000). Phosphorylation of Syk activation loop tyrosines is essential for Syk function. An in vivo study using a specific anti-Syk activation loop phosphotyrosine antibody. *J Biol Chem*, 275, 35442-7. ↗

### Editions

2012-01-04	Authored, Edited	Garapati, P V.
2013-02-13	Reviewed	Roncagalli, R.