

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 70

This document contains 1 pathway and 3 reactions ([see Table of Contents](#))

SMG1 phosphorylates UPF1 (enhanced by Exon Junction Complex) ↗


Location: Nonsense Mediated Decay (NMD) enhanced by the Exon Junction Complex (EJC)

Stable identifier: R-SPO-927889

Type: transition

Compartments: cytosol

Inferred from: SMG1 phosphorylates UPF1 (enhanced by Exon Junction Complex) (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

SMG6 hydrolyzes mRNA with premature termination codon ↗


Location: Nonsense Mediated Decay (NMD) enhanced by the Exon Junction Complex (EJC)

Stable identifier: R-SPO-927836

Type: omitted

Compartments: cytosol

Inferred from: SMG6 hydrolyzes mRNA with premature termination codon (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: [Decay of mRNA in SMG6:SMG5:SMG7:mRNA complex](#)

Decay of mRNA in SMG6:SMG5:SMG7:mRNA complex ↗


Location: Nonsense Mediated Decay (NMD) enhanced by the Exon Junction Complex (EJC)

Stable identifier: R-SPO-927830

Type: omitted

Compartments: cytosol

Inferred from: Decay of mRNA in SMG6:SMG5:SMG7:mRNA complex (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](/electronic_inference_compara.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: SMG6 hydrolyzes mRNA with premature termination codon

Table of Contents

Introduction	1
☒ Nonsense Mediated Decay (NMD) enhanced by the Exon Junction Complex (EJC)	2
☞ SMG1 phosphorylates UPF1 (enhanced by Exon Junction Complex)	3
☞ SMG6 hydrolyzes mRNA with premature termination codon	4
☞ Decay of mRNA in SMG6:SMG5:SMG7:mRNA complex	5
Table of Contents	6